



Autonomic Testing Preparation Instructions

Autonomic testing, also known as autonomic reflex screen or autonomic response testing, is a non-invasive test that measures how the nervous system works to control blood pressure, heart rate and sweating. Your doctor or neurologist may recommend autonomic testing if you're experiencing symptoms including:

- Abnormal sweating
- Dizziness
- Fainting
- Fatigue
- Fluctuating blood pressure
- Numbness and burning in your hands or feet
- Rapid heart rate
- Shortness of breath

Your doctor may order this test for other reasons as well.

Autonomic testing results

Physiological measurements will be obtained in both parts of the test (the QSART and the cardiovascular autonomic test with head-up tilt) to determine evidence of dysautonomia. The results can help physicians diagnose, evaluate, monitor and treat disorders of the autonomic nervous system, including:

- Abnormal sweating
- Autonomic neuropathy
- Baroreflex failure
- Certain degenerative diseases, such as multiple system atrophy and autonomic failure in Parkinson's disease
- Diabetes
- Hypotension
- Orthostatic intolerance
- Orthostatic cerebral hypoperfusion
- Postural tachycardia
- Pure autonomic failure
- Supine hypertension
- Syncopal seizures
- Syncope

What to expect during autonomic testing:

Autonomic testing consists of activities designed to stimulate the autonomic nervous system to produce changes in blood pressure, heart rate, breathing and sweating in a controlled setting. The test includes two main parts - a quantitative sudomotor axon reflex test (if ordered) and a cardiovascular autonomic test.

- **Quantitative sudomotor axon reflex test (QSART)**

QSART is a test that measures the nerves that control sweating. During this first part of the autonomic test, we apply a mild electrical current to your skin to activate the nerves that supply the sweat glands. You may experience mild burning or tingling sensations in the areas being stimulated.

- **Cardiovascular autonomic test with head-up tilt**

During this second part of the autonomic test, we record and monitor your blood pressure, respiration, heart rate, and heart rhythm. This test involves performing simple maneuvers such as taking deep breaths, blowing into a mouth piece and changing position from lying down to standing upright. This test involves performing the simple maneuvers described below.

1. Deep breathing

This test examines how the heart rate responds to breathing slowly and deeply for a minute and 20 seconds at a rate of six breaths per minute.

2. Valsalva maneuver

This test involves blowing forcefully into a mouthpiece against a closed airway. The Valsalva maneuver examines how the body compensates for changes in the amount of blood that returns to the heart.

3. Head-up tilt

You must be laying down for at least 45 minutes prior to starting this test so this test will be done last. This test evaluates your blood pressure, heart rate, and breathing rate in response to standing up. You will start by lying on a stretcher-like table, relaxing and breathing normally, with straps attached around your abdomen and legs for. After 5 minutes of laying flat we will then tilt the table up to a 70-degree angle for 10 minutes to reproduce your symptoms in a controlled setting while continuously monitoring your blood pressure, heart rate, and breathing rate. If you experience symptoms - such as lightheadedness, weakness or feeling faint (known as pre-syncope) — we will tilt the table back to the lying position.

Preparation: (VERY IMPORTANT) - Autonomic testing takes 1.5 to 2 hours.

Please arrive 15 minutes early, if you are late or not properly prepared for testing, it may be necessary to reschedule your test. In the event you are unable to keep your appointment, please call us at (760) 631-3000 to cancel your appointment. We ask that you cancel at least 24 hours in advance.

- The listed medications attached are medications that can affect Autonomic Studies and may hinder you from getting accurate results of your testing. Not all medications are listed and need to be discussed with your ordering provider before this test can be scheduled. All drugs that can affect your sweat, heart rate and blood pressure should be addressed by the ordering provider. We cannot give you advice as to the safety of discontinuing your medications. **These medications must be verified with the prescribing provider's approval and instructions before stopping.** The number of days or doses needed to be stopped prior to testing is indicated in the column next to the medication name. **If these medications are not stopped at the time of your appointment you may be asked to reschedule your appointment.** If any of the listed medications cannot be stopped, please notify the provider that ordered this test before continuing with the Autonomic Testing as this may prevent getting accurate results and they may tell you not to move forward with this test. There may also be other medications not listed that the ordering provider has requested you to hold, if so please follow your provider's instructions and let the tech know when you arrive to your appointment. Please also note, all over the counter (OTC) cold and/or allergy medications should be stopped 3 days prior to testing unless otherwise noted. See attached form. If no medication instruction list is attached you may also obtain it by going online to the following website or please contact our office:

https://neurocenter.com/diagnostic_testing/autonomic-testing/

- Wear comfortable, loose-fitted clothing (loose pants/sweats or shorts and a loose t-shirt). No wired or tight fitting bra's. Do not wear double layered clothing. If your referring provider ordered this test with sweat test (QSART) please be aware that during the testing, one pant leg will need to be raised above the knee. You will be asked to remove your shoes and socks.
- Shower the morning of your appointment or the night before.
- No Topicals (oils or lotions) on your skin the day of test.
- Shave your forearms and legs from (foot to knee only) the day before. This applies to all patients who are being tested for QSART (sweat test) only.
- No nicotine, alcohol, marijuana or caffeine use Twenty-Four (24) hours before the test.
- Do not eat anything heavy 3 hours before the test, you can have a light meal.
- Morning of the appointment no corsets, compressions stockings or anything of that nature 24 hrs prior.
- Please make sure you are well hydrated the day before your test. Please note you will not be able to get up or use the restroom during testing.

You will be laying down flat for approximately an hour to an hour and a half and then at the end tilted up to a standing position for 10 min during this test. Please note you will also not be able to get up at all during this test. If you do not think you're are capable of doing so please contact us.

Risks of autonomic testing

Autonomic testing is considered safe. Some people may faint or collapse during the head-up tilt. When a patient's symptoms and continuous recording of heart rate and blood pressure suggest pre-syncope (near fainting), the patient will be tilted back to a flat position immediately.

After the test: When your test is complete, you can return to your normal activities. Fatigue and lightheadedness may occur, if that pertains to you, you may consider arranging for a friend or family member to pick you up after testing. Please note it takes up to 2 weeks for these results to be interpreted. If there are any urgent finding you will be contacted immediately otherwise your results will be given at you next follow up appointment.

Please call our office if you need to reschedule this appointment. There is a \$200 no show fee if you cancel within 24 hours or if you do not show up. Our phone number is (760) 631-3000.

Thank you!

Autonomic Testing Medication Instructions: The listed medications below are medications that can affect Autonomic Studies and may hinder you from getting accurate results of your testing. Not all medications are listed and need to be discussed with your ordering provider before this test can be scheduled. All drugs that can affect your sweat, heart rate and blood pressure should be addressed by the ordering provider. We cannot give you advice as to the safety of discontinuing your medications. **These medications must be verified with the prescribing provider's approval and instructions before stopping.** The number of days or doses needed to be stopped prior to testing is indicated in the column next to the medication name. **If these medications are not stopped at the time of your appointment you may be asked to reschedule your appointment.** If any of the listed medications cannot be stopped, please notify the provider that ordered this test before continuing with the Autonomic Testing as this may prevent getting accurate results and they may tell you not to move forward with this test. There may also be other medications not listed that the ordering provider has requested you to hold, if so please follow your provider's instructions and let the tech know when you arrive to your appointment. Also note all over the counter cold and allergy medications should be stopped 3 days prior to testing.

Medication:	●	Medication:	●	Medication:	●	Medication:	●	Medication:	●	Medication:	●
Acebutolol	1 day	Clomipramine	7 days	Hytrin	3 days	Orvaten	1 day	Tagamet	1 day		
Adalat	1 day	Clonidine	5 days	Imdur	1 day	Oxybutinin	2 days	Tamsulosin	3 days		
Adderall	3 days	Cogentin	2 days	Imodium	5 days	Pamelor	7 days	Tapentadol	1 day		
Albuterol	1 day	Compazine	5 days	Indapamide	4 days	Paroxetine	7 days	Tenormin	5 days		
Aldomet	1 day	Concerta	3 days	Inderal	2 days	Paxil	7 days	Terbutaline	1 day		
Allegra	3 days	Coreg	2 days	Ivabradine	5 days	Pepcid	1 day	Theo-dur	1 day		
Alprazolam	1 day	Corgard	5 days	Labetalol	3 days	Periactin	2 days	Theophylline	1 day		
Alupent	1 day	Corlanor	5 days	Lavacol	2 days	Phenergan	2 days	Tiazac	2 days		
Amantidine	7 days	Cyclobenzaprine	7 days	Lexapro	7 days	Phentolamine	3 days	Thorazine	7 days		
Aminophylline	1 day	Cymbalta	7 days	Levsin	2 days	Pilocarpine	1 day	Toprol	2 days		
Amitriptyline	7 days	Cyproheptadine	2 days	Loperamide	2 days	Pindolol	2 days	Toprol XL	5 days		
Amlodipine	4 days	Desipramine	7 days	Lopressor	5 days	Pramipexole	7 days	Trandate	2 days		
Anafranil	7 days	Desyrel	7 days	Loratidine	3 days	Prazosin	1 day	Trazodone	7 days		
Antivert	2 days	Detrol	2 days	Lotrel	4 days	ProAmatine	1 day	Trental	1 day		
Apresoline	2 days	Dicyclomine	2 days	Lozide	4 days	Procardia	1 day	Trihexyphenidyl	1 day		
Artane	3 days	Diltiazem	2 days	Luvox	5 days	Prochlorperazine	5 days	Trinalin	3 days		
Atarax	3 days	Dimenhydrinate	3 days	Meclizine	2 days	Promethazine	2 days	Valsartan	2 days		
Atenolol	5 days	Dimetapp	2 days	Mestinon	1 day	Propranolol	2 days	Venlafaxine	5 days		
Atomoxetine	7 days	Diovan	2 days	Metaproterenol	2 days	Propulsid	1 day	Ventolin	2 days		
Atrovent	2 days	Diphenhydramine	2 days	Methyl dopa	1 day	Provigil	3 days	Verapamil	2 days		
Aventyl	7 days	Dipivefrin	3 days	Methylphenidate	3 days	Prozac	7 days	Vicodin	1 day		
Axid	1 day	Ditropan	2 days	Metoclopramide	3 days	Pseudoephedrine HCl	2 days	Visken	2 days		
Azatadine	3 days	Donnatal	5 days	Metoprolol Succinate	5 days	Pyridostigmine	1 day	Vistaril	2 days		
Benadryl	2 days	Doxazosin	5 days	Metoprolol Tartrate	2 days	Quetiapine	2 days	Wellbutrin XL/SR	5 days		
Benicar	3 days	Doxepin	7 days	Midodrine	1 day	Ranitidine	1 day	Xanax	1 day		
Bentyl	2 days	Dramamine	3 days	Minipress	1 day	Regitine	1 day	Zantac	1 day		
Benylin	2 days	Duloxetine	7 days	Mirabegron	5 days	Reglan	3 days	Zebeta	3 days		
Benzotropine	2 days	Effexor	5 days	Mirapex	7 days	Reserpine	7 days	Zofran	1 day		
Betapace	4 days	Elavil	7 days	Modafinil	3 days	Resperidone	5 days	Zoloft	7 days		
Bisoprolol	3 days	Escitalopram	7 days	Morphine IR/SR	<u>5 doses</u>	Risperdal	5 days	Zyrtec	2 days		
Bupropion XL/SR	5 days	Excederin	1 day	Myrbetriq	5 days	Ritalin	1 day				
Buspar	1 day	Famotidine	1 day	Nadolol	5 days	Ritalin LA	3 days	*OTC cold/allergy	3 days		
Buspiron	1 day	Fexofenadine	3 days	Nebivolol	4 days	Robinul	1 day				
Bystolic	4 days	Flexeril	7 days	Nefazodone	7 days	Salagen	1 day				
Calan	2 days	Flomax	3 days	Nifedipine	3 days	Salmeterol	1 day				
Cardizem	2 days	Florinef	5 days	Norco	1 day	Scopolamine	3 days				
Cardura	2 days	Fludrocortisone	5 days	Normadyne	2 days	Sectral	1 day				
Carvedilol	2 days	Fluoxetine	7 days	Norpramin	7 days	Serevent	4 days				
Catapres	5 days	Fluvoxamine	5 days	Nortriptyline	7 days	Seroquel	2 days				
Celexa	7 days	Glycopyrrolate	2 days	Norvasc	4 days	Sertraline	5 days				
Cetirizine	2 days	Guanfacine	5 days	Nucynta	1 day	Serzone	7 days				
Chlorpromazine	7 days	Hydralazine	2 days	Olanzapine	3 days	Sotalol	4 days				
Cimetidine	1 day	Hydrocodone/Acetaminophen	1 day	Olmesartan	3 days	Strattera	7 days				
Cisapride	2 days	Hydroxyzine	3 days	Olopatadine HCL	2 days	Sudafed	2 days				
Citalopram	7 days	Hyoscyamine	2 days	Ondansetron	1 day	Symmetrel	7 days				
Claritin	2 days	Hyperstat	2 days	Orphenadrine	3 days	Synacort	1 day				

