

## VNG Testing Medications Instructions



Certain medications can influence the body's response to the test, which can provide false or misleading results.

In preparation for the VNG test, please review the list below. **DO NOT TAKE them 48 hours prior to your test.**

Please discuss with your prescribing physician if you can safely hold them.

Transdermal Scopolamine patch must remain off for 72 hours prior to testing

Additionally, in the 48 hours prior to the test, DO NOT consume caffeine of any kind, alcohol, nicotine, or marijuana.

Aleve	Bronkotabs	Cymbalta	Dyazide	lasix	Milltown	Oxycontin	Pseudoephedrine	Surax	Ultracet
Abilify	Bucladine	Cystospaz	Effexor	Levsin	Mitran	Pamelor	Ranitidine	Surmontil	Ultram
Actifed	Bucladlin-S Softabs	Dallergy Cap	Elavil	Lexapro	Moban	Paraflex	Remeron	Tagamet	Unisom
Adderall	Bucizine HCL	Dalmane	Endep	L-Hyoscyamine	Morphine	Parafon Forte	Restoril	Talacen	Valium
Allegra	Buspar	Dalmane	Endocet	Librax	Morphine Sulfate	Patanase	Ritalin	Tanxene	Vertigoheel
Ambien	Buspirone	Darvocet	Entex IA	Libritabs	Motrin	Paxil	RMS	Tavist	Vicodin
Amitriptyline	Carbamate	Darvon	Equinil	Librium	MS Contin	Paxipam	Robaxin	Tedrol	Vistaril
Amoxapine	Carisoprodol	Deconamine	Estazolam	Limbitrol	MSIR	Pediacare	Robaxin	Tegretol	Vivactil
Antihistamine Sprays	Celexa	Decongestant	Ethotoin	Lioresal	Mudrane	Peganone	Roxanol SR	Temaril	Vontrol
Antivert	Centrax	Demerol	Etrafon	Lorazepam	Mysoline	Pentobarbital	Roxicodone	Temazepam	Vyvanse
Asendin	Cephalon	Desyrel	Extendryl	Loricet	Naproxen Sodium	Pepcid	Rozerem	Thorazine	Wellbutrin
Astenin Spray	Chlordiazepoxide	Dexedrine	Fioricet	Lortab	Nembural	Percadan	Ru-vert	Tigan	Wygesic
Astepro Spray	Chlorephenesin	Dextrostat	Flexeril	Lovelco	Neptazone	Percocet	Rynatan	Tofranil	Xanax
Atarax	Chlort-trimiten	Diazepam	Floriset	Lufyllin-EPG	Neurontin	Percodone	Salsalate	Topical	Zanaflex
Ataraz	Chlorzoxazone	Dilantin	Flurazepam	Lunesta	Norco	Pergaset	Scopolamine	Tramadol	Zantac
Atavert	Clarinet	Dilaudid	Forflex	Maolate	Norflex	Phenaphen w/ Codeine	Secobarbital	Transderm	Zofran
Ativan	Claritin	Dimhydrinate	Fortis	Marezine	Norgesic	Phenergan	Seconal	Tranxene	Zolof
Baclofen	Clemastine	Dimetabs	Gabapentin	Maxide	Norpramin	Phenobarbital	Seldane	Trazodone	Zyrtec
Barbidonna	Clonazepam	Diphenhydramine	Gemonil	Mebaral	Novafed	Phenylpropamine	Serax	Triaminic	
Belladonna Alkaloids	Clorazepate	Diphenicior	Halazepam	Meclizine	Novahistine	Phenytoin	Serzone	Triavil	
Belladonna	Clozaril	Dipostassium	Halcion	Mepergan	Nyquil	Prazepam	Silenor	Triazolam	
Bellafoline	Codeine	Dixmiss	Haldol	Meperidine HCl	O-Flex	Prilosec	Sinequan	Trimeprazine	
Bellergal-S Tab	Codimal-A	Dolene	Hismanol	Mephentoin	Orflagen	Primidone	Skelaxin	Trimethobenzamide HLC	
Benedryl	Compazine	Donnatal	Hydrocodone	Mephobarbital	Ornade	Promethazine	Sodol	Trinal	
Benylin	Concerta	Doral	Hydromorphone	Merezin	Orphenadizine	Propoxyphine	Soma	Trinxene	
Benzadioepine	Cumetidine	Dorcol	Hydroxyzine	Mesantoin	Ouadrinal	Propoxyphene	Sonata	Tussend	
Bonine	Cyclizene	Dramamine	Imipromaine	Methabital	Oucizepam	Prosum	Stadol	Tylenol PM	
Bromfed	Cyclobenzaprine	Drixoral	Kadain	Methadone	Oxazepam	Provigil	Stelatine	Tylenol w/Codeine	
Bronkoxir	Cylert	Durgesic	Klonopin	Methocarbamol	Oxycodone	Prozac	Sudafed	Tylox	